

## Impact of storms along the coastline of Emilia-Romagna: the morphological signature on the Ravenna coastline (Italy)

P. Ciavola†, C. Armaroli†, J. Chiggiato‡, A. Valentini‡, M. Deserti‡, L. Perini∞ and P. Luciani∞

† Dip. Scienze della Terra, Università di Ferrara, via Saragat 1, 44100 Ferrara, Italy  
Email: cvp@unife.it

‡ ARPA-SIM, Viale Silvani 6 and 2/3, 40122 Bologna, Italy

∞ Servizio Geologico, Sismico e dei Suoli, Direzione Generale Ambiente e Difesa del Suolo e della Costa, Regione Emilia-Romagna, Viale Silvani 4/3, 40122 Bologna, Italy



### ABSTRACT

P. CIAVOLA, C. ARMAROLI, J. CHIGGIATO, A. VALENTINI, M. DESERTI, L. PERINI and P. LUCIANI, 2007. Impact of storms along the coastline of Emilia-Romagna: the morphological signature on the Ravenna coastline (Italy). Journal of Coastal Research, SI 50 (Proceedings of the 9th International Coastal Symposium), pg – pg. Gold Coast, Australia, ISBN)

The coastal zone of the Emilia-Romagna region is exposed to risk from coastal flooding during storms. The vulnerability of this coastline is a topic of interest for future coastal planning as this area provides large revenue for coastal communities and the whole region. An assessment of the historical record of storm events for the period 2000-2004 was carried out using a wave generation model, fed by archived atmospheric circulation data. High-resolution air-borne laser detection technology (LIDAR) and video-monitoring (ARGUS) have permitted to undertake a feasibility study for the assessment of coastal erosion and flooding patterns along one of the best-preserved dune systems of the area. The source of the work was the comparison between two LIDAR flights at an interval of one year (July 2003-September 2004) with the second flight carried out after a major storm occurred in September 2004, with a return period of 25-years. Following this exceptional event the whole coastal system resulted to be in a state of weakness. The study presented in this paper suggests that for the coast of the Emilia-Romagna region the critical factor that controls dune destruction and inland flooding is the joint occurrence of storm waves and surges. Beach slope is the main morphological control on wave run-up and the impact evaluation for the exceptional event has proved that a wide beach can be considered a symptom of safety. A small beach replenishment was undertaken on the test site but this resulted to be ineffective because of the small volume.

**ADDITIONAL INDEX WORDS:** *LIDAR Surveys, Wave forecasting, Beach recovery, Video imagery*

### INTRODUCTION

Although the study of coastal changes using cartography and aerial photography provides some general clues about large-scale coastal changes, the quantification of processes over time-scales longer than five years becomes difficult, as it involves long-term datasets of waves and morphology which are not always available.

In the case of the quantification of the impact of coastal storms, it is quite rare that pre-storm survey are available, unless routine monitoring is foreseen by competent authorities. In the case of the Emilia-Romagna region, beach profiles are surveyed on average every five years, clearly not enough to describe the impact of extreme events, but not even suited to quantify seasonal changes. Some authors (FERREIRA, 2005; CIAVOLA *et al.*, 2006) have recently concluded that the implication of variability in storminess on coastal morphologies remains badly described in the scientific literature. The main limitation is the availability of representative datasets regarding the morphology. Despite the limited tidal excursion of the Adriatic Sea, much discussion is taking place in Emilia-Romagna about which element should be used to identify erosive beach states (e.g. the shoreline, the dune base, etc.). Following extensive aerial photo-interpretation, CIAVOLA *et al.* (2003) have proposed the use of the High Water Line, which is

however extremely sensitive to variability due to surges and wave set-up. In this sense, the position of the MSL contour, worked out using “virtual” topographic surveys using rectified video images provides a meaningful indicator. As recently pointed out by DAVIDSON *et al.* (2006), video-derived Coastal State Indicators have several advantages over similar parameters obtained using conventional image sources, like air or satellite imagery. On the other hand, video-monitoring only covers small areas. Thus, airplane-based LIDAR survey can augment topographic knowledge and work towards identification of the MSL contour.

The absence of wave data for a particular region can be compensated by the use of wind-driven models. However, the use of wind observations for local forecasting is problematic, due to their punctual nature, while the use of locally calibrated, high-resolution circulation models provides a meaningful source for forecasting. The present paper initially describes the occurrence of high-energy events along the Emilia-Romagna coastline from the records available either as direct measurements or as archived wave forecasts. The paper then studies in detail the impact of storms occurred in the Winter 2004-2005 on a system of coastal dunes south of Ravenna and the associated shoreline changes. Finally, the effectiveness of a replenishment to counteract dune erosion is assessed.

## METHODS

### Wave Data

Buoy observations in Italy are provided by APAT through the RON (Rete Ondametrica Nazionale) network. Along the northern Italian coast of the Adriatic Sea there are two buoys which have collected data in recent years, one located near Ancona (43.83 N; 13.71 E) and one near the Po Delta (44.97 N; 12.63 E). The former has the longest wave record (1999-2006), but one has to remember the distance from the study area (Figure 1).

It was decided to prepare a review on the occurrence of storms over the period September 2000-March 2004. The lack of long-term buoy records from the upper Adriatic made impossible to evaluate these extreme events from observations. It was therefore decided to use wave forecasts from the WAM models used by ARPA-SIM to characterise the sea state as a part of their statutory duties for the local government. For this purpose, model runs were extracted at a point with geographical coordinates of 44.5 N and 12.33 E, ideally located a water depth of 9 m just south of Ravenna.

The spectral Wave Model Cycle4 (WAMC4) developed by the WAMDI GROUP (1988) was used for wave prediction. The considered domain was the whole Adriatic Sea area, with a 8 km horizontal resolution ( $1/12^\circ$ ). WAM was forced using the wind computed at 10 m above sea level by the meteorological model LAMBO. As described by MARSIGLI *et al.* (2001), the Limited Area Model Bologna (LAMBO) was the limited area numerical weather prediction model that had been operational until March 2004 at ARPA-SIM. This was a grid-point, split-explicit, primitive equation model, based on an early version of the NCEP Eta Model (MESINGER *et al.*, 1988). The prognostic variables were zonal and meridional wind components, specific humidity, temperature and surface pressure. LAMBO was operationally integrated twice a day, nested on ECMWF operational runs of 00 and 12 UTC, the forecast length being 72 and 84 hours, respectively. Verification of WAM wave forecasts against observations at Ancona indicate that it was possible to use the [+24,+48 hours] forecast data provided by the WAM model for the isolation and evaluation of extreme events like sea storms. For further details can be found in the final report of the CADSEALAND Project (2006).

Many methods can be applied to isolate an extreme storm event from a set of samples, each method differing from the others for the considered variables and their combination (e.g. MENDOZA and JIMÉNEZ, 2004). For the identification and isolation of sea storms it has been considered as threshold a significant wave height higher than 1.5 m and a minimum event duration of 6 hours both for forecast data and observations.

### Field Site and Morphological Datasets

The case study site (Lido di Dante) is characterized by a beach stretching 3 km from the edge of coastal protection structures to the Bevano river mouth (Figure 1). In front of the urbanised area there is a breakwater (770 m long) and its edge effects generated by the southern end of the breakwater are felt as far as 900 m from the structures, especially during NE storms (ARMAROLI *et al.*, 2005). The area south of the village is instead free from coastal structures, with a pine forest and an eroding dune field. The dune system is single-ridged, with crest elevation of 1-1.5 m above MSL on the northern part and up to 5 m above MSL at the southern margin. On February 2003 an ARGUS video-system was installed at the site (Figure 1). The system consists of four cameras mounted on a 18 m-high tower: three looking at the beach behind the breakwater, and one looking at the natural sector. In




Figure 1. Location of Lido di Dante, Ravenna (© CGR Parma, AIMA flight 1996).

this study the shoreline was extracted using the IBM (Intertidal Beach Mapper) tool of AARNINKHOF *et al.* (2003). Short-term (hours) shoreline surveys were carried out by ARMAROLI *et al.* (2006) to compare the DGPS data with ARGUS using a GIS. The study revealed that ARGUS and DGPS shorelines are identified with comparable precision.

Two topographic LIDAR flights were undertaken at an interval of approximately one year (July 2003-September 2004). The first flight was provided courtesy of ENI, while the second one was carried out by the SGSS of the Emilia-Romagna region. The second flight took place after a major storm, occurred on 24 September 2004. The morphology of the beach and the dunes was also monitored in 2004/2005 along cross-shore profiles regularly spaced (150 m), starting from the dune crest down to a depth of -1/-1.5 m below MSL. The dune crest elevation was also measured walking on the foredune ridge with an RTK-DGPS in dynamic mode. Particular attention was put on measuring the extension of overwash areas and their elevation.

## RESULTS AND DISCUSSION

### Storm Climatology

Because of space constraints, only the time series of the WAM archive and Ancona buoy are presented here. The five year WAM dataset shows that storms mostly occur in the autumn and winter seasons (Figure 2). The main direction of incoming waves during storms is the North-East in Winter and the East in Autumn. The frequency of occurrence seems to have grown during the winter period of the last three considered years, whereas it has remained more or less constant for the autumn period. Springs storms are less pronounced both in frequency of occurrence and strength, but they show less directional variability than autumn events, being all concentrated around the North-East.

These results are partially confirmed by the analysis of buoys observations, taking into account the distance of Ancona and Punta della Maestra buoys from the Emilia Romagna coastline and the different water depths below the buoys compared to the extrapolated point of the regional domain where simulation has been carried out.




Figure 2. Seasonal number and direction of sea storm forecasted by WAM model or observed by the Ancona buoy.

The Ancona buoy (depth of 75 m) has a long and consistent record of data, but it is also further away from the domain of interest. The other buoy of Punta della Maestra (depth of 30 m) is closer, but on the other hand it has a short and discontinuous dataset. For this reason in Figure 2 only a comparison with between WAM and Ancona is presented. To notice that the forecast database only covered the period up to the Spring of 2004. Thereafter a major model re-organizations took place and for consistency was not considered further use.

**The Storms of September 2004**

Between 18:30 GMT of 24 September and 00:30 GMT of 27 September the buoy at Ancona registered an  $H_s$  of more than 5.5 m. If one uses a storm threshold for the wave height of 1.5 m, this was exceeded for 73 hours between 24th and 27th September

(Figure 3). During the event there was also a maximum storm surge of 0.7 m measured by the tide gauge of Porto Corsini, near Ravenna. The value of the surge was computed comparing the astronomical tide prediction with the tide gauge record. The surge was not particularly high, below the level of the 1-yr event indicated in YU *et al.* (1998). Offshore wave data identify a first storm occurring between 24-25 September and a second one between 26-27 September. The maximum energy was reached during the first event. Transposition of the wave offshore climate to the study site predicts a wave height of 5.65 m, which exceeded the 25-yr return event.

In Figure 3 it is possible to notice that the maximum water level registered by the tide gauge in Ravenna did not correspond to the storm peak. Wave direction changed during the storm from SE to NNE. The SE winds caused the surge because they were combined with low barometric pressure. The effect of the storm on the coast could have been worse if the surge and the maximum tidal level reached on 24 September 2004 were simultaneous with the storm peak.

**The Morphological Record of Storminess**

From the analyses of storminess presented above, it is clear that the Winter of 2003/2004 was an exceptional year for what concerns the number of storms from NE, with good agreement between the wave forecasts and the buoys. On the other hand, the Autumn of 2004 was characterised by the concomitant action of exceptional high tides and storm clustering from the North. At this stage it was decided to investigate two issues. First to test the response of the beach at Lido di Dante following this “exceptional” clustering of events that occurred in September 2004. Secondly, to assess if there was a trend of continuous erosion that was onset by the frequent storms of the previous years.

To evaluate the erosion caused by storms in Autumn 2004 and whether or not the beach recovered its equilibrium state, the shoreline evolution was assessed from video images. Choosing days with spring tides under calm conditions, images of the area immediately before the storm were compared with others captured immediately after the event and at the following spring tides.

The results (Figure 4) show that the area is divided into two parts: the one close to the structures was dramatically eroding, the central and southern areas were oscillating or stable.

The video images also showed that the dunes were touched by run-up and overtopped by waves. Regarding the central and southern parts of the beach, the analysis of video images confirmed that were rhythmic features moving alongshore,




Figure 3. A- Significant wave height ( $H_s$ ) and wave directions (Dir) registered by the Ancona buoy between 23 and 28 September 2004; B- Residuals between measured tides at Porto Corsini (Ravenna) and astronomical predictions.




Figure 4. Comparison between MSL identified using ARGUS imagery on 2 September 2004 and on 30 September 2004. The interruption in the interpretation is due to the presence of a large bush which covers part of the beach. Notice also the oscillation in the shoreline due to formation migration of swash bars.

according to the dominant wave direction, as proposed by ARMAROLI *et al.* (2005).

The rhythmic features are strictly related to the formation and migration of intertidal bars, which are common at the site (BALOUIN *et al.*, 2006). The analysis of the shoreline position revealed that the southern part of the beach is wide and “safe”. Regarding the dune foot, retreat in the central area was observed, while in the area close to the structures the dunes almost disappeared.

The analysis of the topographic datasets collected during 2003/2004 proved that an acceleration in the erosion rate of the dunes was caused by overwash and overtopping events, that however had started during storms in the Winter of 2003/2004. A comparison between the dune profiles measured in the surveys before and after the storm occurred in September 2004 (Figure 5), revealed a variable behaviour of the area up to about 1 km from the structures, where it was actually achieved a peak in retreat (14 m), during that single event. Profiles showed that there were localised sediment inputs due to dune erosion during the storms, with many points of overwashing observed in the field.

It is striking how the dune undercutting rate dropped passed the 1 km boundary from the structures, which also corresponded to a general stability of the intertidal beach observed from the video imagery and the topographic surveys. To notice that this area receives consistent sediment input from lateral erosion of dunes at the Bevano river mouth, due to northward migration of the inlet and subsequent sand removal from the mouth by long-shore drift directed northwards (CIAVOLA *et al.*, 2005).

The dunes that suffered most from overtopping were those close to the structures in an area that historically suffers from consistent beach erosion (ARMAROLI *et al.*, 2005). In Figure 5, it is possible

to notice the strong retreat of the beach and the destruction of the first line of dunes. It is important to notice how the areas with healthy dunes (over 4 m high) were not affected at all by the exceptional event of September 2004. Where the dune system is stable and the beach slope mild the beach tended to be wider.

A conceptual model of the morphological behaviour of this beach when hit by storms was produced dividing the area into three zones. A first one located between the edge of the groin and a boundary at -350 m to the south, with high erosion rates. The issue to be considered in this case is the influence of the breakwater. When north-easterlies generate storm waves, diffraction by the barrier’s edge may generate a concentration of wave energy on this coastal stretch. The beach slope was indeed generally steep in the area immediately to the south of the barrier. A second intermediate coastal segment, extended up to -1050 m southwards of the tower, had variable erosion/accretion tendencies. On the other hand, the southernmost area, which extended to the edge of the coverage by ARGUS (-1200 m), was instead essentially stable, even under extreme forcing conditions, with combined occurrence of high waves and surge levels.

### Beach Recovery Potential

A replenishment scheme was carried out in the Spring of 2005 with the placement of 33000 m<sup>3</sup> along the beach south of the structures, along a distance of 800 m from the breakwater.

To assess the effectiveness of the replenishment, two DGPS surveys were undertaken before the beginning of the works (27 February 2005) and after completion (7 May 2005). The survey consisted of 20 profiles closely spaced every 50 m (Figure 6), extended from the dune crest to the low tide limit. The same profiles were then extracted from the DTM of the LIDAR Survey of SGSS undertaken in September 2004, after the storms that eroded the dunes. The aims of the study was to evaluate the evolution of the beach during the Winter, the effectiveness of the replenishment and the displacement of the sediment dumped during operations. As it can be seen in Figure 6, the beach had not recovered from the erosive trend that was onset by the storms of September 2004 and just few weeks after replenishment most of the sand had been removed from the dumping site (northern profiles, e.g. n. 20).

It seemed however that the sand was displaced southward and re-arranged on the intertidal beach in the form of swash/bars and low tide terraces, which are typical of the area. A volumetric comparison between the surveys before and after the works confirm a discontinuous pattern of erosion/deposition around mean sea level. Clearly the sand volume injected into the littoral budget was not large enough to produce tangible results.




Figure 5. Comparison between 2003-2004 LIDAR data of two profiles that are immediately to the south from the protected area. Notice the destruction of the first line of dunes.




Figure 6. Comparison between beach profiles surveyed after major storm (Sept 04), winter 2005 and after beach replenishment (May 05).

## CONCLUSIONS

The frequency of occurrence of storm has grown during the winter period of the last three years considered (2002-2004), whereas it has been more or less constant for the autumn period. The maximum number of storms occurred during the 2003-2004 winter period. During September 2004 a group of strong sea storms caused hazards for open sea and harbour activities, damage to seaside tourist facilities and severe erosion of the Emilia-Romagna littoral. The work on the case study site of Lido di Dante has quantified the impact of this exceptional storm, with combined high waves and storm surge, on one of the few remaining areas in the Region with coastal dunes.

The storms generated beach loss mainly in the area close to the structures. Here the time sequence of the storms played an important role. The erosive effect was enhanced by the joint occurrence of storms in a short time, the so-called "clustering effect" (FERREIRA, 2005). This led to the incapacity of the system to recover an equilibrium state between two sequences of "wave attacks". Emergency replenishment schemes were not successful, only providing a "placebo" effect on the sediment budget.

## ACKNOWLEDGEMENTS

The research was financed by the INTERREG IIIB-CADSES project CADSEALAND, partially supported by the EU. Installation of the ARGUS system was sponsored by the EU project COASTVIEW (contract EVK3-CT-2001-00054). We are grateful to S. Caleffi and M. Gardelli for support during data analysis.

## LITERATURE CITED

- AARNINKHOF, S.G.J., TURNER, I.L., DRONKERS, D.T., CALIJOUW, M. and NIPIUS, L., 2003. A video based technique for mapping intertidal beach bathymetry. *Coastal Engineering*, 49, 275-289.
- ARMAROLI C., BALOUIN Y., CIAVOLA P. and CAPATTI D., 2005. Nearshore bars as a natural protection of beaches, field evidences from Lido di Dante beach, Adriatic Sea. *Proceedings of ICCCM'05 Conference*, University of Porto, Portugal, pp. 295-303.
- ARMAROLI, C., CIAVOLA, P., BALOUIN, Y. and GATTI, M., 2006. An integrated study of shoreline variability using GIS and ARGUS techniques. *Journal of Coastal Research*, SI 39, 473-477.
- BALOUIN, Y., CIAVOLA, P. and ARMAROLI, C., 2006. Sediment transport patterns and coastal evolution at Lido di Dante beach, Adriatic Sea. *Proceedings of Coastal Dynamics 2005*, ASCE, New York, USA.
- CADSEALAND Project, 2006. Analysis of correlation between coast evolution and meteo-marine climatology. Deliverable Report 4.5, Available at: [http://www.regione.emilia-romagna.it/wcm/geologia/appuntamenti/09\\_11\\_06\\_cadsealand/Rapporto\\_Finale\\_CADSEALAND.pdf](http://www.regione.emilia-romagna.it/wcm/geologia/appuntamenti/09_11_06_cadsealand/Rapporto_Finale_CADSEALAND.pdf)
- CIAVOLA, P., CORBAU, C., CIBIN, U. and PERINI, L., 2003. Mapping of the coastal zone of the Emilia-Romagna region using Geographical Information Systems. *Proceedings of MEDCOAST 2003 Conference*, E. OZHAN (ed.), MEDCOAST, Ravenna (Italy), pp. 2363-2374.
- CIAVOLA P., BILLI P., ARMAROLI C., PRECISO E., SALEMI E. and BALOUIN Y., 2005. Valutazione della morfodinamica di foce del Torrente Bevano (RA): il ruolo del trasporto solido di fondo. *Geologia Tecnica ed Ambientale*, 1/2005, 41-57
- DAVIDSON, M.A., AARNINKHOF, S.G.J., VAN KONINGSVELD, M. and HOLMAN, R.A., 2006. Developing coastal video monitoring systems in support of coastal zone management. *Journal of Coastal Research*, SI 39, ICS 2004, Brazil.
- FERREIRA, O., 2005. Storm groups versus extreme single storms: predicted erosion and management consequences. *Journal of Coastal Research*, SI 42, 221-227.
- MARSIGLI C., MONTANI A., NEROZZI F., PACCAGNELLA T, TIBALDI S., MOLteni F. and BUIZZA R., 2001. A strategy for high-resolution ensemble prediction. II: Limited-area experiments in four Alpine flood events. *Q. J. R. Meteorol. Soc.*, 127, 2095-2115.
- MENDOZA, E.T. and JIMENEZ J.A., 2004. Factors controlling vulnerability to storm impacts along the Catalan coast. *Proceedings International Conference of Coastal Engineering 2004*, Lisbon, pp. 3087-3099.
- MESINGER F., JANJIC Z. I., NICKOVIC S., GAVRILOV D. and DEAVEN D. G., 1988. The step-mountain co-ordinate: Model description and performance for cases of Alpine lee cyclogenesis and for a case of Appalachian redevelopment. *Mon. Wea. Rev.*, 116, 1493-1518.
- YU C.S., DECOUTTERE C. and BERLAMONT J. 1998. Storm surge simulations in the Adriatic Sea. *CENAS*, Kluwer Academic, Dordrecht, The Netherlands, pp. 207-232.